Author index

Ahlers, J., see Arnold, W.M. (942) 83
Ahlers, J., see Arnold, W.M. (942) 96
Alonso, A., see Valpuesta, J.M. (942) 341
Arnold, W.M., Zimmermann, U., Heiden, W. and Ahlers, J.
The influence of tetraphenylborates (hydrophobic anions) on yeast cell electro-rotation (942) 96
Arnold, W.M., Zimmermann, U., Pauli, W., Benzing, M., Niehrs, C. and Ahlers, J.
The comparative influence of substituted phenols (especially chlorophenols) on yeast cells assayed by electro-rotation and other methods (942) 83
Arrondo, J.L.R., see Valpuesta, J.M. (942) 341

Barisas, B.G., see Rosenthal, K.S. (942) 38
Basset, G., Saumon, G., Bouchonnet, F. and Crone, C.
Apical sodium-sugar transport in pulmonary epithelium in situ (942) 11
Bennekou, P. and Stampe, P.
The effect of ATP, intracellular calcium and the anion exchange inhibitor DIDS on conductive anion fluxes across the human red cell membrane (942) 179
Benzing, M., see Arnold, W.M. (942) 83
Bevers, E.M., see Verhallen, P.F.J. (942) 150
Bouchonnet, F., see Basset, G. (942) 11
Brzezinski, P., Malmström, B.G., Lorentzon, P. and Wallmark, B.
The catalytic mechanism of gastric H^+/K^+-ATPase: simulations of pre-steady-state and steady-state kinetic results (942) 215

Chang, C.L., see Chang, T.-M. (942) 57
Chang, T.-M. and Chang, C.L.
Hepatic uptake of asialoglycoprotein is different among mammalian species due to different receptor distribution (942) 57
Cheeseman, C.I., see Lee, C.W. (942) 139
Chifu, E., see Tomoaia-Cotisel, M. (942) 295
Comfurius, P., see Verhallen, P.F.J. (942) 150
Cox, T.C.
Potassium dependence of sodium transport in frog skin (942) 169
Crone, C., see Basset, G. (942) 11

De Kruijff, B., see Zhou, L.-X. (942) 115
Devaux, P.F., see Rosso, J. (942) 271
Dhers, C., see Leonards, K.S. (942) 245
Dobrović-Jenik, D. and Milković, S.
Regulation of fetal Na^+/K^+-ATPase in rat kidney by corticosteroids (942) 227

Favilli, F., see Vincenzini, M.T. (942) 107
Fernández, M.S.
^{1}H-NMR of phosphatidylcholine liposomes at low p^2H in the presence of a paramagnetic shift reagent (942) 199
Flores, E., see Schmetterer, G. (942) 33
Freysz, L., see Pelletier, X. (942) 125

Giraldez, F., Valverde, M.A. and Sepúlveda, F.V.
Hypotonicity increases apical membrane Cl^- conductance in *Necturus* enterocytes (942) 353
Goñi, F.M., see Valpuesta, J.M. (942) 341
Grossman, R.A., see Hoshiko, T. (942) 186

Harris, W.E. and Stahl, W.L.
Origin of the γ polypeptide of the Na^+/K^+-ATPase (942) 236
Heiden, W., see Arnold, W.M. (942) 96
Hertz, E., see Streichman, S. (942) 333
Hoshiko, T., Grossman, R.A. and Machlup, S.
Effects of basolateral ouabain, amphotericin B, cyanide and potassium on amiloride noise during voltage clamp of *Rana pipiens* skin support sodium-amiloride competition (942) 186

Iantomasi, T., see Vincenzini, M.T. (942) 107
Ise, N., see Kitano, H. (942) 131

Jarvis, S.M., see Lee, C.W. (942) 139
Jessen, H., Vorum, H., Jørgensen, K.E. and Sheikh, M.I.
Characteristics of D-alanine transport by luminal membrane vesicles from pars convoluta and pars recta of rabbit proximal tubule (942) 262
Jordi, W., see Zhou, L.-X. (942) 115
Jørgensen, K.E., see Jessen, H. (942) 262

Kato, N., see Kitano, H. (942) 131
Kitano, H., Kato, N., Tanaka, N. and Ise, N.
Mutual recognition between polymerized liposomes: enzyme and enzyme inhibitor system (942) 131
Kneer, C., see Stevens, B.R. (942) 205
Koynova, R.D., see Quinn, P.J. (942) 315

Lee, C.W., Cheeseman, C.I. and Jarvis, S.M.
Na^+- and K^+-dependent uridine transport in rat renal brush-border membrane vesicles (942) 139
Le Grimellec, C., see Venien, C. (942) 159
Leonards, K.S. and Dhers, C.
Changes in the surface charge properties of isolated cardiac sarcolemmal vesicles measured by light scattering. II. Characteristics of rabbit preparations (942) 245

Leray, C., see Pelletier, X. (942) 125
Lis, L.J., see Quinn, P.J. (942) 315
Lis, L.J., see Tenchov, B.G. (942) 305
Lorentzon, P., see Brzezinski, P. (942) 215
Lucas, M., Mata, R. and Romero, A.
Comparison of the active calcium extrusion, calcium buffering capacity and ATPase activity in rabbit reticulocytes and mature red cells (942) 65

Macarulla, J.M., see Valpuesta, J.M. (942) 341
Machlup, S., see Hoshiko, T. (942) 186
Malmström, B.G., see Brzezinski, P. (942) 215
Mata, R., see Lucas, M. (942) 65
McMurchie, E.J. and Patten, G.S.
Dietary cholesterol influences cardiac β-adrenergic receptor adenylate cyclase activity in the marmoset monkey by changes in membrane cholesterol status (942) 324
Mellor, I.R., Thomas, D.H. and Sansom, M.S.P.
Properties of ion channels formed by *Staphylococcus aureus* δ-toxin (942) 280
Milković, S., see Dobrović-Jenik, D. (942) 227
Mocanu, A., see Tomoaia-Cotisel, M. (942) 295

Nagle, J.F. and Wiener, M.C.
Structure of fully hydrated bilayer dispersions (942) 1
Nagy, K.
Conditional open and delay time histograms of sodium channels (942) 209
Niehrs, C., see Arnold, W.M. (942) 83

Ojcius, D.M. and Solomon, A.K.
Sites of *p*-chloromercuribenzenesulfonate inhibition of red cell urea and water transport (942) 73
Ortonne, J.-P., see Verrando, P. (942) 45

Patten, G.S., see McMurchie, E.J. (942) 324
Pauli, W., see Arnold, W.M. (942) 83
Pelletier, X., Freysz, L. and Leray, C.
Topological distribution of choline phospholipid fatty acids in trout intestinal brush-border membrane (942) 125
Pimplikar, S.W. and Reithmeier, R.A.F.
Studies on the interaction of matrix-bound inhibitor with Band 3, the anion transport protein of human erythrocyte membranes (942) 253
Pisani, A., see Verrando, P. (942) 45

Quinn, P.J., Koynova, R.D., Lis, L.J. and Tenchov, B.G.
Lamellar gel–lamellar liquid crystal phase transition of dipalmitoylphosphatidylcholine multilayers freeze-dried from aqueous trehalose solutions. A real-time X-ray diffraction study (942) 315
Quinn, P.J., see Tenchov, B.G. (942) 305
Quinn, P.J., see Tomoaia-Cotisel, M. (942) 295

Reithmeier, R.A.F., see Pimplikar, S.W. (942) 253
Reithmeier, R.A.F., see Werner, P.K. (942) 19
Roess, D., see Rosenthal, K.S. (942) 38
Romero, A., see Lucas, M. (942) 65
Rosenthal, K.S., Roess, D. and Barisas, B.G.
Herpes simplex virus type 1 penetration initiates mobilization of cell surface proteins (942) 38
Rosso, J., Zachowski, A. and Devaux, P.F.
Influence of chlorpromazine on the transverse mobility of phospholipids in the human erythrocyte membrane: relation to shape changes (942) 271

Sansom, M.S.P., see Mellor, I.R. (942) 280
Saumon, G., see Basset, G. (942) 11
Schmetterer, G. and Flores, E.
Uptake of fructose by the cyanobacterium *Nostoc* sp. ATCC 29150 (942) 33
Sepúlveda, F.V., see Giraldez, F. (942) 353
Sheikh, M.I., see Jessen, H. (942) 262
Solomon, A.K., see Ojcius, D.M. (942) 73
Stahl, W.L., see Harris, W.E. (942) 236
Stampe, P., see Bennekou, P. (942) 179
Stevens, B.R. and Kneer, C.
Lanthanide-stimulated glucose and proline transport across rabbit intestinal brush-border membranes (942) 205
Streichman, S., Hertz, E. and Tatarsky, I.
Direct involvement of spectrin thiols in maintaining erythrocyte membrane thermal stability and spectrin dimer self-association (942) 333

Tanaka, N., see Kitano, H. (942) 131
Tatarsky, I., see Streichman, S. (942) 333
Tenchov, B.G., Lis, L.J. and Quinn, P.J.
Structural rearrangements during crystal–liquid-crystal and gel–liquid-crystal phase transitions in aqueous dispersions of dipalmitoylphosphatidylethanolamine. A time-resolved X-ray diffraction study (942) 305
Tenchov, B.G., see Quinn, P.J. (942) 315
Thomas, D.H., see Mellor, I.R. (942) 280
Tomoaia-Cotisel, M., Zsako, J., Mocanu, A., Chifu, E. and Quinn, P.J.
Monolayer properties of membrane lipids of the extreme halophile *Halobacterium cutirubrum*, at the air/water interface (942) 295
Tsai, A.-l., Vijjeswarapu, H. and Wu, K.K.
Interaction between platelet receptor and iloprost isomers (942) 220

Valpuesta, J.M., Goñi, F.M., Alonso, A., Arrondo, J.L.R. and Macarulla, J.M.
Lipid–protein interactions. The mitochondrial complex III-phosphatidylcholine-water system (942) 341
Valverde, M.A., see Giraldez, F. (942) 353
Venien, C. and Le Grimellec, C.
Phospholipid asymmetry in renal brush-border membranes (942) 159
Verhallen, P.F.J., Bevers, E.M., Comfurius, P. and Zwaal, R.F.A.
Fluoride-dependent calcium-induced platelet procoagulant activity shows that calpain is involved in increased phospholipid transbilayer movement (942) 150

Verrando, P., Pisani, A. and Ortonne, J.-P.
The new basement membrane antigen recognized by the monoclonal antibody GB3 is a large size glycoprotein: modulation of its expression by retinoic acid (942) 45
Vijjeswarapu, H., see Tsai, A.-l. (942) 220
Vincenzini, M.T., Favilli, F. and Iantomasi, T.
Glutathione-mediated transport across intestinal brush-border membranes (942) 107
Vorum, H., see Jessen, H. (942) 262

Wallmark, B., see Brzezinski, P. (942) 215
Werner, P.K. and Reithmeier, R.A.F.
The mechanisms of inhibition of anion exchange in human erythrocytes by 1-ethyl-3-[3-(trimethylammonio)propyl]carbodiimide (942) 19
Wiener, M.C., see Nagle, J.F. (942) 1
Wu, K.K., see Tsai, A.-l. (942) 220

Zachowski, A., see Rosso, J. (942) 271
Zhou, L.-X., Jordi, W. and De Kruijff, B.
Influence of heme and importance of the N-terminal part of the protein and physical state of model membranes for the apocytochrome *c*–lipid interaction (942) 115
Zimmermann, U., see Arnold, W.M. (942) 83
Zimmermann, U., see Arnold, W.M. (942) 96
Zsako, J., see Tomoaia-Cotisel, M. (942) 295
Zwaal, R.F.A., see Verhallen, P.F.J. (942) 150